


Background of the Invention

The present invention relates to a novel process for the preparation of (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene of the formulae



or salts thereof, or the D- or L-hydrogentartrates thereof and also their further conversion to give (1S,4R)- or (1R,4S)-4-(2-amino-6-chloro-9-H-purine-9-yl)-2-cyclopentene. (1R,4S)-1-Amino-4-(hydroxymethyl)-2-cyclopentene of the formula IV is an important intermediate for the preparation of carbocyclic nucleosides such as, for example, Carbovir® (Campbell et al., J. Org. Chem. 1995, 60, 4602-4616).

A process for the preparation of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene is described, for example, by Campbell et al. (ibid) and by Park K.H. & Rapoport H. (J. Org. Chem. 1994, 59, 394 - 399). In this process, the starting material is either D-glucono-δ-lactone or D-serine, approximately 15 synthesis stages being required to form (1R,4S)-N-tert-butoxycarbonyl-4-hydroxymethyl-2-cyclopentene, and the protecting group is removed to give (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene.

Both these processes are costly, complex and not practicable industrially. WO 93/17020 describes a process for the preparation of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene, in which (1R,4S)-4-amino-2-cyclopentene-1-carboxylic acid is reduced to the desired product using lithium aluminium hydride.

Disadvantages of this process are firstly that the double bond of the cyclopentene ring is also reduced, the poor handling properties of lithium aluminium hydride and secondly that it is too costly.

Taylor S.J. et al. (Tetrahedron: Asymmetry Vol. 4, No. 6, 1993, 1117 - 1128) describe a process for the preparation of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene


019842 112498
86427 2486160

starting from (\pm)-2-azabicyclo[2.2.1]hept-5-en-3-one as starting material. In this process, the starting material is converted, using microorganisms of the species *Pseudomonas solanacearum* or *Pseudomonas fluorescens*, into
5 (1R,4S)-2-azabicyclo[2.2.1]hept-5-en-3-one, which is then reacted with di-tert-butyl dicarbonate to give (1R,4S)-N-tert-butoxycarbonyl-2-azabicyclo[2.2.1]hept-5-en-3-one, and the latter is reduced using sodium borohydride and trifluoroacetic acid to give the desired product.

10 This process is far too costly.
In addition, Martinez et al. (J. Org. Chem. 1996, 61, 7963 - 7966) describe a 10-stage synthesis of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene starting from diethyl dialkylmalonate. This process too has the disadvantage that it is complex and not practicable industrially.
15

It is also known that N-substituted (\pm)-2-azabicyclo[2.2.1]hept-5-en-3-ones, which carry an electron-withdrawing substituent, can be reduced to the corresponding
20 N-substituted aminoalcohols using a metal hydride (Katagiri et al., Tetrahedron Letters, 1989, 30, 1645 - 1648; Taylor et al., *ibid*).

In contrast to this, it is known that unsubstituted (\pm)-2-azabicyclo[2.2.1]hept-5-en-3-one of the
25 formula



II

is reduced with lithium aluminium hydride to give (\pm)-2-azabicyclo[2.2.2]octene (Malpass & Tweedle, J. Chem. Soc., Perkin Trans 1, 1977, 874 - 884), and that the direct reduction of (\pm)-2-azabicyclo[2.2.2]hept-5-en-3-one to give the corresponding aminoalcohol has to date
30 been impossible (Katagiri et al., *ibid*; Taylor et al., *ibid*).

It is also known to resolve racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene using (-)-dibenzoyl-tartaric acid (US-A 5 034 394). On the one hand, this
35

09198427 112498


a

reaction has the disadvantage that (-)-dibenzoyltartaric acid is expensive, and, on the other hand, that the separation must take place in the presence of an exactly defined mixture of acetonitrile and ethanol. This solvent mixture cannot be removed and must be fed to the combustion.

Summary of the Invention


The object of the present invention was to provide a simple, economical and cost-effective process for the preparation of a (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene.

Surprisingly, it has now been found that when (+)-2-azabicyclo[2.2.1]hept-5-en-3-one of the formula



II

in the form of the racemate or one of its optically active isomers, is reduced with a metal hydride, the aminoalcohol of the formula



I

in the form of the racemate or one of its optically active isomers is obtained in a simple manner.

Detailed Description

As the person skilled in the art is aware, the aminoalcohol of the formula I can be converted using an acid into the corresponding salts, such as, for example, into hydrohalide salts. Suitable hydrohalide salts are hydrobromides and hydrochlorides.

The starting material, the (+)-2-azabicyclo[2.2.1]hept-5-en-3-one can be prepared according to EP-A 0 508 352.

Metal hydrides which may be used are alkali metal or alkaline earth metal hydrides and also binary or complex metal hydrides of the boron or aluminium group, such as alkali metal and alkaline earth metal borohydrides, alkali metal and alkaline earth metal aluminium hydrides. Suitable alkali metal or alkaline earth metal

hydrides are LiH, NaH, KH, BeH₂, MgH₂ or CaH₂.

Binary alkali metal or alkaline earth metal borohydrides which may be used are NaBH₄, LiBH₄, KBH₄, NaAlH₄, LiAlH₄, KAlH₄, Mg(BH₄)₂, Ca(BH₄)₂, Mg(AlH₄)₂ and Ca(AlH₄)₂. Complex metal hydrides of the boron or aluminum group may have the general formula M¹M²H_nL_m, in which n is an integer from 1 to 4, and m is an integer from 4 to 4 minus the corresponding number n, M¹ is an alkali metal atom, M² is boron or aluminium, and L is C₁₋₄-alkyl, C₁₋₄-alkenyl, C₁₋₄-alkoxy, CN or an amine, or the complex metal hydrides may have the general formula M²H_oL_p, in which M² is as defined above and o is an integer from 0 to 3, and p is an integer from 3 to 3 minus the corresponding number p. Possible M¹M²H_nL_m compounds are LiBH(C₂H₅)₃, LiBH_x(OCH₃)_{4-x}, LiAlH(OC(CH₃)₃)₃, NaAlH₂(OC₂H₄OCH₃)₂, NaAlH₂(C₂H₅)₂ or NaBH₃CN. Preferably, the reduction is carried out using a metal borohydride. As an expert in the art is aware, the metal hydrides mentioned such as, for example, LiBH₄, can also be produced "in situ". Common preparation methods for LiBH₄ are, for example, the reaction of an alkali metal borohydride with a lithium halide (H.C. Brown et al., Inorg. Chem. 20, 1981, 4456 - 4457), the reaction of LiH with B₂O₃ in the presence of hydrogen and a hydrogenation catalyst (EP-A 0 512 895), the reaction of LiH with (H₅C₂)OBF₃ (DE-A 94 77 02) and that of LiH with B(OCH₃)₃ (US-A 2,534,533).

The metal hydrides are expediently used in a molar ratio of from 1 to 5 per mole of (±)-2-azabicyclo-[2.2.1]hept-5-en-3-one.

The metal hydrides, in particular NaBH₄, are preferably used with lithium salt additives. Lithium salts which may be used are LiCl, LiF, LiBr, LiI, Li₂SO₄, LiHSO₄, Li₂CO₃, Li(OCH₃) and LiCO₃.

The reduction is expediently carried out in an inert-gas atmosphere, such as, for example, in an argon or nitrogen atmosphere.

The reduction can be carried out at a temperature of from -20 to 200°C, preferably at a temperature of from

SECRET 112498 a

60 to 150°C.

Suitable solvents are aprotic or protic organic solvents. Suitable aprotic organic solvents may be ethers or glycol ethers, such as, for example, diethyl ether, dibutyl ether, ethyl methyl ether, diisopropyl ether, tert-butyl methyl ether, anisole, dioxane, tetrahydrofuran, monoglyme, diglyme and formaldehyde dimethylacetal. Suitable protic organic solvents are C₁₋₆-alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, tert-butanol, pentanol, tert-amyl alcohol or hexanol and also mixtures of these with water. Suitable protic organic solvents are also mixtures of one of said ethers, glycol ether with water or with one of said alcohols, such as a mixture of a C₁₋₆-alcohol with an ether or glycol ether, in particular a mixture of methanol, ethanol or water with diethyl ether, tetrahydrofuran, dioxane, glyme or diglyme. The solvent used is preferably a protic organic one, such as a mixture of a C₁₋₆-alcohol or water with an ether or glycol ether.

In a preferred embodiment, the reduction is carried out in the presence of an additive such as in the presence of water or of a ^{univalent or polyvalent} ~~lower aliphatic~~ alcohol. The ^{univalent C₁₋₆} ~~lower aliphatic~~ alcohol ^{can be chosen from} may be methanol, ethanol, methoxyethanol, n-propanol, isopropanol, isobutanol, tert-butanol, n-butanol, ^{and} ~~The polyvalent C₁₋₆ alcohol can be chosen from~~ diols such as butanediol, and triols such as glycerol. In particular, the ~~lower aliphatic~~ alcohol is methanol or ethanol. Here, the ~~lower aliphatic~~ alcohol is expediently used in a molar ratio of from 2 to 15 per mol of (+)-2-azabicyclo[2.2.1]hept-5-en-3-one.

If the reaction is carried out in the presence of said alcohol, the corresponding amino acid ester can be formed in situ (intermediate). I.e. if the starting material used is (+)-2-azabicyclo[2.2.1]hept-5-en-3-one, according to the invention the corresponding (+)-amino acid ester can be formed. If the starting material used is (-)-2-azabicyclo[2.2.1]hept-5-en-3-one, according to the invention the (-)-amino acid ester can correspondingly be formed as intermediate.


09198427 112498

a
a


a
a
a
a

its substituent like
Katzgrub does.
Intermediate
etc

Surprisingly, it has also been found that when a cyclopentene derivative of the general formula



in the form of the racemate or one of its optically active isomers, in which R is C₁₋₄-alkyl, C₁₋₄-alkoxy, aryl or aryloxy, is hydrolyzed with an alkali metal hydroxide, the aminoalcohol of the formula



in the form of the racemate or one of its optically active isomers is obtained in a simple manner.

C₁₋₄-Alkyl can be substituted or unsubstituted. In the text below substituted C₁₋₄-alkyl is taken to mean C₁₋₄-alkyl substituted by ^{one or more} halogen atoms. The halogen atom may be F, Cl, Br or I. Examples of C₁₋₄-alkyl are methyl, ethyl, propyl, butyl, isobutyl, tert-butyl, isopropyl, chloromethyl, bromomethyl, dichloromethyl and dibromomethyl. The C₁₋₄-alkyl is preferably methyl, ethyl, propyl, butyl, isobutyl or chloromethyl.

The C₁₋₄-alkoxy used may be, for example, methoxy, ethoxy, propoxy or butoxy. The aryl used can be, for example, phenyl or benzyl, substituted or unsubstituted.


The aryloxy used can be, for example, benzyloxy or phenoxy, substituted or unsubstituted.

The alkali metal hydroxide used may be sodium or potassium hydroxide.

For this process variant, the cyclopentene derivative of the general formula III is preferably prepared by reduction of the corresponding acyl-2-azabicyclo[2.2.1]hept-5-en-3-one of the general formula

09198427 112498

Ind.
a²



IV


in the form of the racemate or one of its optically active isomers, in which R is as defined above, using one of the metal hydrides already mentioned in an anhydrous solvent.

5 The anhydrous solvent may be protic or aprotic organic solvents, in particular an anhydrous protic organic solvent such as a tertiary alcohol. The tertiary alcohol may be tert-butyl alcohol or tert-amyl alcohol.


10 As already mentioned above, this reduction is also preferably carried out in the presence of an addition, such as in the presence of a ^{C1-6}~~lower~~ aliphatic alcohol such as methanol, in particular in the presence of 2 mol of methanol per mole of acyl-2-azabicyclo-[2.2.1]hept-5-en-3-one (formula IV).

15 The reaction is expediently carried out at a temperature of from 0 to 50°C, preferably from 15 to 30°C.

20 The racemic aminoalcohol, ^{preferably the cis-racemic,} of the formula I is then converted according to the invention either by chemical means using an optically active tartaric acid or by biotechnological means using a hydrolase in the presence of an acylating agent to give (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene of the formula




V




VI

25 or salts thereof and/or to give (1S,4R)- or (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene derivative of the general formulae



VII



VIII

or salts thereof, in which X and Y are identical or different and are an acyl group or H, with the exception of X = Y = H.

The hydrolases used may be lipases, proteases, amidases or esterases, lipases being expediently used.

In the text below, salts are taken to mean hydrohalide salts such as hydrochlorides, hydrobromides or tartrates.

As the person skilled in the art is aware, hydrolase-catalyzed acylations in which optically active compounds are formed are carried out in the presence of a suitable acylating agent (Balkenhohl et al., 1997, J. Prakt. Chem. 339, 381 - 384; K. Faber, "Biotransformation in Organic Chemistry", 2nd ed., Berlin 1995, 270 - 305). Suitable acylating agents are generally carboxylic acid derivatives such as carboxamides, carboxylic anhydrides or carboxylic esters. The carboxylic esters may, for example, be alkoxycarboxylic esters, such as ethyl methoxyacetate and isopropyl methoxyacetate, C₁₋₆-carboxylic esters, such as butyl acetate, ethyl butyrate and ethyl hexanoate, glyceryl esters, such as tributyrin (glyceryl tributyrate), glycol esters, such as glycol dibutyrate and diethyl diglycolate, dicarboxylic esters, such as diethyl fumarate and malonate, cyanocarboxylic esters, such as ethyl cyanoacetate, or cyclic esters, such as, for example, 6-caprolactone.

Accordingly, the acyl group in the formulae VII and VIII corresponds to the acid component of the carboxylic acid derivative used.

The lipases used may be standard commercial lipases, such as, for example: Novo lipase SP523 from *Aspergillus oryzae* (Novozym 398), Novo lipase SP524 from *Aspergillus oryzae* (lipase = Palatase 20000 L from Novo), Novo lipase SP525 from *Candida antarctica* (lipase B Novozym 435, immobilized), Novo lipase SP526 from *Candida*

09158427 112498

antarctica (lipase A = Novozym 735, immobilized), lipase
kits from Fluka (1 & 2), Amano P lipase, lipase from
Pseudomonas sp., lipase from Candida cylindracea, lipase
from Candida lipolytica, lipase from Mucor miehei, lipase
5 from Aspergillus niger, lipase from Bacillus thermo-
catenulatus, lipase from Candida antarctica, lipase AH
(Amano; immobilized), lipase P (Nagase), lipase AY from
Candida rugosa, lipase G (Amano 50), lipase F (Amano F-
AP15), lipase PS (Amano), lipase AH (Amano), lipase D
10 (Amano), lipase AK from Pseudomonas fluorescens, lipase
PS from Pseudomonas cepacia, newlase I from Rhizopus
niveus, lipase PS-CI (immobilized lipase from Pseudomonas
cepacia). These lipases may, as the person skilled in the
art is aware, be used as cell-free enzyme extracts or
15 else in the corresponding microorganism cell.

The proteases may also be commercially available,
such as, for example, serine proteases such as sub-
tilisins. The subtilisin may be savinase from Bacillus
sp., alcalase, subtilisin from Bacillus licheniformis and
20 also proteases from Aspergillus, Rhizopus, Streptomyces
or Bacillus sp.

The biotechnological racemate resolution is expediently
carried out at a temperature of from 10 to 80°C and at a
pH of from 4 to 9.

25 The biotechnological racemate resolution is
expediently carried out in a protic or aprotic organic
solvent. Suitable aprotic organic solvents are ethers
such as tert-butyl methyl ether, diisopropyl ether,
dibutyl ether, dioxane and tetrahydrofuran, aliphatic
30 hydrocarbons such as hexane, organic bases such as
pyridine, and carboxylic esters such as ethyl acetate,
and suitable protic organic solvents are the C₁₋₆-alcohols
already described, such as, for example, pentanol.

The (1S,4R)- or (1R,4S)-1-amino-4-(hydroxy-
35 methyl)-2-cyclopentene derivatives of the general for-
mulae VII and VIII formed in accordance with the inven-
tion during the biotechnological racemate resolution are,
depending on the desired target compound (aminoalcohol of
the formula V or VI), hydrolyzed by chemical means to

09198427-112498

give the aminoalcohol of the formula V or VI. The chemical hydrolysis is expediently carried out in an aqueous basic solution or using a basic ion exchanger. The aqueous basic solution is preferably, as for the hydrolysis of the cyclopentene derivatives of the general formula III described above, an alkali metal hydroxide. The basic ion exchangers can, for example, be Dowex 1x8(OH⁻) and Duolite A147.

The chemical racemate resolution is carried out using an optically active tartaric acid such as using D-(-)-tartaric acid or L-(+)-tartaric acid.

The racemate resolution with D-(-)-tartaric acid is expediently carried out by firstly reacting the racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene with the D-(-)-tartaric acid in the presence of a ^{C1-6}~~lower aliphatic~~ alcohol.

Suitable ^{C1-6}~~lower aliphatic~~ alcohols are the same as those described above. Preference is given to using methanol. The reaction which leads to formation of the salt is usually carried out at temperature between 20°C and the reflux temperature of the solvent, preferably at the reflux temperature.

If desired, the 1-amino-4-(hydroxymethyl)-2-cyclopentene D-tartrate formed during the reaction can be further purified by recrystallization from ^{the C1-6}~~a lower aliphatic~~ alcohol such as methanol.

The racemate resolution with L-(+)-tartaric acid is expediently carried out as that with D-(-)-tartaric acid. I.e. the racemate resolution with L-(+)-tartaric acid is likewise carried out in the presence of a lower aliphatic alcohol and at a temperature between 20°C and the reflux temperature of the solvent, preferably at the reflux temperature. After cooling, the (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene L-hydrogentartrate crystallizes out.

The (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene L-hydrogentartrate is present, in particular, in dissolved form in the mother liquor.

Isolation, further purification (liberation) and

09198427-112498

a

a

a

conversion to the corresponding salt of (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene takes place with a base and subsequent acid treatment. Suitable bases are alkali metal alkoxides, alkali metal or alkaline earth metal carbonates, or alkali metal or alkaline earth metal hydroxides. The alkali metal alkoxides may be sodium or potassium alkoxides. The alkali metal carbonate may be potassium or sodium carbonate, potassium or sodium hydrogencarbonate, and the alkaline earth metal carbonate may be magnesium or calcium carbonate. The alkali metal hydroxide may be sodium or potassium hydroxide, and the alkaline earth metal hydroxide may be calcium hydroxide. Conversion to the corresponding salt usually takes place with a mineral acid such as with sulphuric acid, hydrochloric acid or phosphoric acid, preferably with hydrochloric acid.


(1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene D-hydrogentartrate and (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene L-hydrogentartrate are compounds unknown in the literature and are likewise provided by the invention.

Preference is given to carrying out the chemical racemate resolution with D-(+)-tartaric acid due to the higher performance, technical facility and more efficient racemate resolution.

As for the racemic aminoalcohol, it is of course also possible to react the optically active (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentenenes with D-(-)- or L-(+)-tartaric acid to give the corresponding tartrates.

A further constituent of the present invention is the further conversion, the acylation, of the (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentenenes to give the (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene derivative of the general formula

09198427.112498




Here, the substituent R is as defined in the cyclopenten derivative of the general formula III.

The acylation can be carried out using a carbonyl halide of the general formula



- 5 in which X is a halogen atom, and R is as defined above, or using a carboxylic anhydride of the general formula



in which R is as defined above.

The halogen atom X may be F, Cl, Br or I. Preference is given to Cl or F.

- 10 Examples of carbonyl halides are: acetyl chloride, chloroacetyl chloride, butyryl chloride, isobutyryl chloride, phenylacetyl chloride, benzyl chloroformate, propionyl chloride, benzoyl chloride, alkyl chloroformate or tert-butyloxycarbonyl fluoride.

- 15 Examples of carboxylic anhydrides are: tert-butoxycarbonyl anhydride, butyric anhydride, acetic anhydride or propionic anhydride. The acylation is preferably carried out using a carboxylic anhydride, in particular using tert-butoxycarbonyl anhydride.



- 20 The acylation can be carried out without solvent or using an aprotic organic solvent. The acylation is expediently carried out in an aprotic organic solvent. Suitable aprotic organic solvents are, for example, pyridine, acetonitrile, dimethylformamide, diisopropyl ether, tetrahydrofuran, toluene, methylene chloride, 25 N-methylpyrrolidone, triethylamine, chloroform, ethyl acetate, acetic anhydride and mixtures thereof.

The acylation is expediently carried out at a


09198427 112498

temperature of from -20 to 100°C, preferably from 0 to 80°C.



The further conversion according to the invention of (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene D- or L-hydrogentartrate to (1S,4R)- or (1R,4S)-4-(2-amino-6-chloro-9-H-purine-9-yl)-2-cyclopentenyl-1-methanol, or a salt thereof, of the formulae



is carried out by reacting (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene D- or L-hydrogentartrate with N-(2-amino-4,6-dichloropyrimidin-5-yl)formamide of the formula



to give (1S,4R)- or (1R,4S)-4-[(2-amino-6-chloro-5-formamido-4-pyrimidinyl)amino]-2-cyclopentenyl-1-methanol of the formulae



and then cyclizing the latter in a known manner to give the compounds according to formula ~~VIII~~ and ~~IX~~.

N-(2-Amino-4,6-dichloropyrimidin-5-yl)formamide can be prepared according to WO 95/21 161.

The reaction is expediently carried out in the presence of a base. Suitable bases are the same as those previously described for liberating (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentenenes from the corresponding tartrate.


The reaction is expediently carried out in a protic solvent. The protic solvent may be ~~lower aliphatic~~ ^{C₁₋₆} alcohols such as methanol, ethanol, propanol, isopropanol, butanol or isobutanol.

The (1S,4R)- or (1R,4S)-4-[(2-amino-6-chloro-5-formamido-4-pyrimidinyl)amino]-2-cyclopentenyl-1-methanol of the formula ~~XIV~~ ^{XI} or ~~XV~~ ^{XII} is then cyclized in a known manner according to WO 95/21 161 to give the end product according to Formula ~~VIII~~ ^{XI} or ~~IX~~ ^{XII}.


The cyclization is usually carried out dissolved in trialkyl orthoformate in the presence of a concentrated aqueous acid. The trialkyl orthoformates used may be, for example, trimethyl or triethyl orthoformate.

The aqueous acid may be, for example, hydrogen fluoride, sulphuric acid or methanesulphonic acid.


A further constituent of the invention is the overall process for the preparation of (1S,4R)-4-(2-amino-6-chloro-9-H-purine-9-yl)-2-cyclopentenyl-1-methanol, or salts thereof, of the formula XII starting from (-)-2-azabicyclo[2.2.1]hept-5-en-3-one or (-)-acyl-2-azabicyclo[2.2.1]hept-5-en-3-one of the formulae




in which R is as defined above, by reduction with a metal hydride to give an aminoalcohol of the formula




or to give a cyclopentene derivative of the general formula




in which R is as defined above, which are then converted into the corresponding hydrohalide salts, and then reacted with N-(2-amino-4,6-dichloropyrimidin-5-yl)-formamide of the formula



- 5 to give (1S,4R)-4-[(2-amino-6-chloro-5-formamido-4-pyrimidinyl)amino]-2-cyclopentenyl-1-methanol of the formula



and then the latter is cyclized in a known manner to give the compound of the formula



- 10 This process variant has the advantage that the hydrohalide salts formed therein may be used as a crude mixture in the preparation of the product of the formula XII.

Examples:

Example 1

Reduction of acyl- or unsubstituted-2-azabicyclo[2.2.1]-hept-5-en-3-one

- a 5 1.1. Preparation of ^{cis}(±)-acetyl-1-amino-4-(hydroxymethyl)-2-cyclopentene in an anhydrous protic organic solvent using sodium borohydride

280 g of 2-methyl-2-butanol (amyl alcohol) and 15.2 g of sodium borohydride (0.4 mol) were charged into a sulphonation flask at 20°C. A mixture of 907 g of (±)-acetyl-2-azabicyclo[2.2.1]hept-5-en-3-one (0.6 mol) and 37.5 g of methanol (2 equivalents based on (±)-acetyl-2-azabicyclo[2.2.1]hept-5-en-3-one was metered into this suspension over the course of 2 h at 20°C. The reaction mixture was then stirred for a further 3 h at 20°C. The solvent was distilled as far as possible (40°C). Boron was removed by adding 280 g of methanol and 27.2 g of formic acid, warming the mixture to 25-30°C and distilling off the methyl borate/methanol azeotrope at this temperature (130 to 80 mbar). The precipitated sodium formate was filtered off, and the filtrate was reduced by evaporation to give 93.4 g of crude product as a clear viscous oil; crude yield: about 84-85%.

- a 1.2. Preparation of [±]cis-1-amino-4-(hydroxymethyl)-2-cyclopentene

A suspension of (±)-2-azabicyclo[2.2.1]hept-5-en-3-one (10.00 g, 91.6 mmol) and lithium borohydride (4.00 g, 183.7 mmol) in dry dioxane (100 ml) was heated in an inert-gas atmosphere (argon) for 4 h at 110°C below the reflux temperature. After this time, about 20-25% of the starting material had reacted to give the product (GC analysis with internal standard benzophenone after work-up of the reaction mixture; work-up: 0.05 ml of the reaction mixture were

quenched with 0.1 ml of 1M HCl and immediately rendered basic using 0.2 ml of 1M NaOH). The structural detection of the product was carried out by H-NMR, GC and GC-MS.

5 1.3. Preparation of cis(+)-1-amino-4-(hydroxymethyl)-2-cyclopentene

10 A 25 ml round-bottom flask was charged with 1.0 g (9.2 mmol) of (+)-2-azabicyclo[2.2.1]hept-5-en-3-one and 0.4 g (18.4 mmol) of lithium borohydride, under an inert-gas atmosphere, in 10 ml of dioxane, and the mixture was refluxed for 3 h at 110°C. Excess reducing agent was destroyed by adding about 5 ml of semi-concentrated HCl (adjusted to pH 3). The mixture was then immediately buffered by adding about 15 1 ml of saturated NaHCO₃ solution at pH 8. GC analysis indicated the formation of the product. The entire reaction mixture was then evaporated to dryness and purified by means of column chromatography (gradient: hexane/ethyl acetate/MeOH = 1:1:1 -> MeOH). In this way cis(+)-2-azabicyclo-20 [2.2.1]hept-5-en-3-one and the corresponding (+)-aminoalcohol were obtained.

25 1.4. Preparation of cis(-)-1-amino-4-(hydroxymethyl)-2-cyclopentene

30 A 25 ml round-bottom flask was charged with 1.0 g (9.2 mmol) of (-)-2-azabicyclo[2.2.1]hept-5-en-3-one and 0.4 g (18.4 mmol) of lithium borohydride, under an inert-gas atmosphere, in 10 ml of dioxane, and the mixture was refluxed for 3 h at 110°C. Excess reducing agent was destroyed by adding about 5 ml of semi-concentrated HCl (adjusted to pH 3). The mixture was then immediately buffered by adding about 35 1 ml of saturated NaHCO₃ solution at pH 8. GC analysis indicated the formation of the product in 18% yield (GC standard is benzophenone). The entire reaction mixture was then evaporated to dryness and

a
5 purified by means of column chromatography (gradient: hexane/ethyl acetate/MeOH = 1:1:1 -> MeOH). In this way, 0.43 g (43%) of ^{cis} \wedge (-)-2-azabicyclo[2.2.1]hept-5-en-3-one was reisolated and 0.04 g (4%) of the corresponding (-)-aminoalcohol was obtained.

By HPLC, only the (-)-enantiomer of the aminoalcohol was detectable. The ee of the product is thus >98%.

a
1.5. Preparation of ^{cis} \wedge (±)-1-amino-4-(hydroxymethyl)-2-cyclopentene in an alcohol

10 A 100 ml round-bottom flask fitted with magnetic stirrer was charged with 3.0 g (27.5 mmol) of (±)-2-azabicyclo[2.2.1]hept-5-en-3-one and 1.2 g (28.3 mmol) of lithium borohydride, under an inert-gas atmosphere, in 35 g of 2-butanol, and the mixture
15 was stirred for 3 h at 60°C. GC analysis of a sample (work-up: 0.1 g sample rendered acidic using 0.2 ml of 1M HCl, then quickly rendered basic using 0.1 ml of saturated NaHCO₃) indicated the formation of the product in 12% yield after this time. (GC standard
20 is benzophenone.)

a
1.6. Preparation of a ^{cis} \wedge (±)-1-amino-4-(hydroxymethyl)-2-cyclopentene in an alcohol/ether mixture

25 A 10 ml round-bottom flask was charged under at inert-gas atmosphere with 0.5 g (4.6 mmol) of (±)-2-azabicyclo[2.2.1]hept-5-en-3-one and 0.59 g (18.4 mmol) of methanol in 7.5 ml of dioxane (abs.). 0.21 g (9.2 mmol) of lithium borohydride were added, and the mixture was heated for 4 h at 60°C. The mixture was then cooled to 5°C using an ice/waterbath, and
30 about 10 ml of semi-concentrated HCl was carefully added to the reaction mixture (vigorous reaction, gas evolution), as a result of which a yellowish clear solution formed. This solution was analyzed directly by a quantitative ion-chromatographic

method. It contained 0.60 mmol (13.1%) of (+)-2-azabicyclo[2.2.1]hept-5-en-3-one (determined as HCl salt of the corresponding amino acid, which is the acidic hydrolysis product of [^]cis(+)-2-azabicyclo[2.2.1]hept-5-en-3-one) and 3.06 mmol of product, corresponding to a yield of 66.8%, aminoalcohol.

1.7. Preparation of [^]cis(+)-1-amino-4-(hydroxymethyl)-2-cyclopentene in the presence of additives such as water or various alcohols

10 A 10 ml round-bottom flask was charged with 0.50 g (4.66 mmol) of (+)-2-azabicyclo[2.2.1]hept-5-en-3-one and 0.30 g (13.7 mmol) of lithium borohydride in 7.5 ml of abs. dioxane, and the mixture was heated to 60°C.

15 At this temperature, over the course of 30 min, X mmol of [^]additive Y such as alcohol or water ~~alcohol~~ was added dropwise using a syringe. The mixture is then stirred for 2 h at 60°C, cooled to about 20°C and poured into about 10 ml of semi-concentrated HCl. The content was then
20 determined directly using a quantitative ion-chromatographic method (cf. Table 1).

09198427 112498
05427 248760

Table 1

Ex-ample	Addi-tive Y	X	X	(±)-2-Aza-bicyclo-[2.2.1]-hept-5-en-3-one	Amino-alcohol
		mmol	equiva-lents	% yield	
1.7.1	-	-	-	15	52
1.7.2	water	17.1	1.25	23.3	67.5
1.7.3	water	34.3	2.5	32.3	58.3
1.7.4	meth-anol	34.3	2.5	4.5	83.1
1.7.5	eth-anol	34.3	2.5	6.5	74.7
1.7.6	isopro-panol	34.3	2.5	28.1	52.3

1.8. Preparation of cis (±)-1-amino-4-(hydroxymethyl)-2-cyclopentene with various amounts of methanol

~~Using the procedure of Example 1.7,~~ the reaction was carried out in a variety of methanol concentrations. The results are given in Table 2.

Table 2

Example	Methanol	Methanol	(±)-2-Azabi-cyclo[2.2.1]-hept-5-en-3-one	Amino-alcohol
	mmol	equivalents	% yield	
1.8.1	9.2	1	27.5	44.8
1.8.2	18.3	2	13.1	66.8
1.8.3	27.5	3	24.7	54.8
1.8.4	36.6	4	5.7	56.8
1.8.5	45.8	5	12.0	58.3
1.8.6	55.0	6	7.2	33.0

1.9 Preparation of ^{cis}(±)-1-amino-4-(hydroxymethyl)-2-cyclopentene with various solvents

~~Using the procedure of Example 1.7, the reaction was carried out in a variety of solvents (7.5 ml) and the content was determined. The results are given in Table 3.~~

5

09198427 112493 054211 22485160

a

ins.
a4

Table 3

Example	Solvent X	(±)-2-Aza-bicyclo-[2.2.1]-hept-5-en-3-one	Amino-alcohol
		% Yield	
1.9.1	dioxane	13.6	79.8
1.9.2	diethyl ether	10.8	68.6
1.9.3	tetrahydrofuran	22.4	67.6
1.9.4	diisopropyl ether	12.6	51.3
1.9.5	tert-butyl methyl ether	10.0	71.3
1.9.6	monoglyme	15.5	75.3
1.9.7	formaldehyde dimethyl acetal	12.0	74.2

1.10 Preparation of $\text{Cis}(\pm)$ -1-amino-4-(hydroxymethyl)-2-cyclopentene with various additions of LiBH_4

~~Following the procedure of Example 1.7,~~ the reaction was carried out using a variety of LiBH_4 concentrations, and the content was determined. The results are given in Table 4.

Table 4

Ex-ample	LiBH ₄	LiBH ₄	(±)-2-Aza-bicyclo-[2.2.1]-hept-5-en-3-one	Amino-alcohol
	mmol	equiva-lents	% yield	
1.10.1	4.6	1	11.9	47.9
1.10.2	6.9	1.5	9.6	45.6
1.10.3	9.2	2	12.7	71.3
1.10.4	11.5	2.5	13.3	74.5
1.10.5	13.8	3	12.8	77.1
1.10.6	16.1	3.5	12.7	62.4

1.11 Preparation ^{cis(+)- or cis(-)}~~(1R,4S) and (1S,4R)~~-1-amino-4-(hydroxy-methyl)-2-cyclopentene in the presence of various alcohols and in the presence of water in a variety of solvents

A 10 ml round-bottom flask fitted with magnetic stirrer was charged with 0.50 g (4.6 mmol) of (±)-2-azabicyclo[2.2.1]hept-5-en-3-one and 0.30 g (13.7 mmol) of lithium borohydride in 6 ml of a variety of solvents, and the mixture was heated to 60°C. At this temperature, over the course of 30 min, 34.3 mmol of the additive Y were added dropwise using a syringe. The mixture was then stirred for 2 h at 60°C, cooled to about 20°C and poured onto about 10 ml of semi-condentrated HCl.

The content is determined directly using a quantitative ion-chromatographic method (cf. Table 5). The ee value of the product was determined by means of HPLC. The results are given Table 5.

Table 5

Example	(-) - 2-Azabicyclo-[2.2.1]hept-5-en-3-one		Solvent	Additive Y	Aminoalcohol	
	(+) - 2-Azabicyclo-[2.2.1]hept-5-en-3-one	ee value			Yield (IC)	ee value (HPLC)
1		98.0	dioxane	water	64.3	>99.0
2		98.0	glyme	water	68.0	>99.0
3		75.9	dioxane	water	65.1	76.0
4		75.9	glyme	water	63.5	75.6
5		50.2	dioxane	water	74.8	51.4
6		51.6	glyme	water	64.1	53.0
7		25.3	dioxane	water	61.1	30.4
8		25.6	glyme	water	61.0	29.6
9		98.0	dioxane	methanol	83.1	98.2
10		98.0	glyme	methanol	81.5	99.2

BOTTLE LABEL

- 25 -

11	75.9		dioxane	methanol	81.4	78.0
12	76.2		glyme	methanol	79.9	78.6
13	50.4		dioxane	methanol	81.3	54.4
14	51.5		glyme	methanol	82.0	55.2
15	24.8		dioxane	methanol	65.2	27.4
16	27.8		glyme	methanol	81.7	32.2
17	98.0		dioxane	ethanol	80.8	80.8
18	98.0		glyme	ethanol	85.1	85.1
19	75.5		dioxane	ethanol	85.3	78.2
20	75.6		glyme	ethanol	83.6	78.4
21	50.7		dioxane	ethanol	76.3	54.4
22	51.1		glyme	ethanol	71.3	55.2
23	25.4		dioxane	ethanol	73.0	28.6
24	25.5		glyme	ethanol	75.0	28.6
25		98.0	dioxane	water	62.0	>99.0

BOHEP-1-CH2F60

26		98.0	glyme	water	59.5	>99.0
27		51.3	dioxane	water	79.0	52.2
28		49.0	glyme	water	61.3	52.0
29		98.0	dioxane	methanol	77.2	>99.0
30		98.0	glyme	methanol	80.0	>99.0
31		49.0	dioxane	methanol	80.8	46.8
32		49.5	glyme	methanol	80.9	48.8

1.12 Preparation of $\text{cis}(\pm)$ -1-amino-4-(hydroxymethyl)-2-cyclopentene using sodium borohydride in various alcohols

Following the procedure of Example 1.7, the reaction was carried out in a variety of ^{additives such as alcohols or water} alcohols. In contrast to Example 1.7, however, sodium borohydride (0.51 g, 13.7 mmol) was used as reducing agent. The results are given in Table 6.

Table 6

Ex-ample	Addi-tive Y	X	X	(\pm)-2-Aza-bicyclo-[2.2.1]-hept-5-en-3-one	Amino-alco-hol
		mmol	equiva- valents	Yield %y	
1.12.1	water	17.1	1.25	75.4	20.1
1.12.2	water	34.3	2.5	71.9	26.7
1.12.3	methanol	34.3	2.5	39.2	22.2
1.12.4	ethanol	34.3	2.5	67.8	8.6
1.12.5	-	-	-	62.2	3.5

1.13 Preparation of $\text{cis}(\pm)$ -1-amino-4-(hydroxymethyl)-2-cyclopentene using NaBH_3CN

60 ml of dioxane and 8.6 g (137 mmol) of sodium cyanoborohydride and 11.9 g (137 mmol) of lithium bromide were refluxed overnight for 15 h at 110°C in a 100 ml sulphonation flask. The mixture was then cooled to 60°C, and a solution of 5.0 g (45.8 mmol) of (\pm)-2-azabicyclo[2.2.1]hept-5-en-3-one containing 15 ml of methanol were added dropwise over the course of 30 min. The white suspension was stirred for 3 h at 60°C, cooled to about 5°C and poured into

about 100 ml of semi-concentrated HCl. The content was then determined directly using a quantitative ion-chromatographic method. The yield of amino-alcohol was about 4%.

5 Example 2

Alkaline hydrolysis of acetyl-(±)-1-amino-4-(hydroxymethyl)-2-cyclopentene

88.9 g of racemic acetyl-1-amino-4-(hydroxymethyl)-2-cyclopentene (content 77.2%) were suspended (partially dissolved) in 70 g of water. 84 g of 30% NaOH (1.1 equivalents) were added thereto, and the solution was refluxed for 3 h. According to TLC, the hydrolysis was complete. The resulting acetate was removed by electrodialysis. The obtained aqueous solution was reduced by evaporation and dried by azeotropic distillation with butanol. The residue was taken up in methanol for racemate resolution. Yield of hydrolysis to (±)-1-amino-4-(hydroxymethyl)-2-cyclopentene was 90%.

20 Example 3: Preparation of (1R,4S)- or (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene

3.1 Racemate resolution using hydrolases

3.1.1 Preparation of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene using lipases

a 25 3.1.1.1 25 mM of racemic ^{cis} 1-amino-4-(hydroxymethyl)-2-cyclopentene were suspended with 1000 units of Novozym 435 in 5 ml of dioxane at room temperature. 25 mM of ethyl methoxyacetate were added as acetylating agent. The formation of N-methoxyacetylaminolcohol was unambiguously detected by TLC. The conversion was 50% (according to estimation of the TLC). This reaction produced (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene.

- 3.1.1.2 50 mM of racemic ^{cis} 1-amino-4-(hydroxymethyl)-2-cyclopentene were suspended with 1000 units (U) of Novozym 435 in 5 ml of tetrahydrofuran. 50 mM of NaOH and 50 mM of ethyl methoxyacetate were added, and the mixture was incubated at 30°C. N-Methoxyacetylalcohol was detected using TLC. The estimated conversion was 50%. This reaction produced (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene.
- 3.1.1.3 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.06 ml of tributyrin (glyceryl tributyrate) and 20 U of Novozym 435 (immob. lipase from Candida antarctica) at room temperature. After 3 days, enantiomerically pure, according to HPLC, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene was obtained in 43% yield.
- 3.1.1.4 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether. 0.02 ml of 6-caprolactone and 20 U of Novozym 435 (immob. lipase from Candida antarctica) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 87% ee was obtained in 49% yield (HPLC).
- 3.1.1.5 100 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of hexane, 0.3 ml of tributyrin and 20 U of Novozym 435 (immob. lipase from Candida antarctica) at room temperature. After 1 week, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 77% ee was obtained in 28% yield (HPLC).

a

09198427 112498

- 3.1.1.6 100 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of tert-butanol, 0.3 ml of tributyrin and 20 U of Novozym 435 (immob. lipase from *Candida antarctica*) at 30°C. After 1 week, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 78% ee was obtained in 15% yield (HPLC).
- 3.1.1.7 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of methyl caproate and 20 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 68% ee was obtained in 52% yield (HPLC).
- 3.1.1.8 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of glycol dibutyrate and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 89% ee was obtained in 31% yield (HPLC).
- 3.1.1.9 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of diethyl fumarate and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 86% ee was obtained in 36% yield (HPLC).
- 3.1.1.10 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of diethyl malonate and 40 U of Novozym 435 (immob. lipase from

05198427 112498

Candida antarctica) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 86% ee was obtained in 21% yield (HPLC).

- 5 3.1.1.11 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of diisopropyl ether, 0.2 mmol of tributyrin and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, enantiomerically pure (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene was obtained in 15% yield (HPLC).
- 10
- 3.1.1.12 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of diisopropyl ether, 0.2 mmol of diethyl fumarate and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 88% ee was obtained in 24% yield (HPLC).
- 15
- 20
- 3.1.1.13 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of diisopropyl ether, 0.2 mmol of diethyl malonate and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 82% ee was obtained in 14% yield (HPLC).
- 25
- 3.1.1.14 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of diisopropyl ether, 0.2 mmol of diethyl diglycolate and 40 U of Novozym 435 (immob. lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 88% ee was obtained in 7%
- 30
- 35

09198427 112458 254277 2486160

yield (HPLC).

- 3.1.1.15 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of dibutyl ether, 0.2 mmol of tributyrin and 40 U of Novozym 435 (immobilized lipase from *Candida antarctica*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 95% ee was obtained in 13% yield (HPLC).
- 3.1.1.16 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of pyridine, 0.02 ml of ethyl 2-methoxyacetate and 20 mg of lipase AK (lipase from *Pseudomonas fluorescens*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 84% ee was obtained in 18% yield (HPLC).
- 3.1.1.17 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of ethyl cyanoacetate and 10 mg of lipase PS (lipase from *Pseudomonas cepacia*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 67% ee was obtained in 40% yield (HPLC).
- 3.1.1.18 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of methyl tert-butyl ether, 0.2 mmol of diethyl fumarate and 10 mg of lipase PS (lipase from *Pseudomonas cepacia*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 86% ee was obtained in 18% yield (HPLC).

0918427-412498
B64277-22486T60

3.1.2 Preparation of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene using proteases

- 3.1.2.1 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of 2-methyl-2-butanol, 0.2 mmol of diethyl maleate and 40 mg of Alcalase (protease from *Bacillus licheniformis*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 28% ee was obtained in 39% yield (HPLC).
- 3.1.2.2 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of 2-methyl-2-butanol, 0.2 mmol of diethyl fumarate and 40 mg of Savinase (protease from *Bacillus* sp.) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 32% ee was obtained in 42% yield (HPLC).
- 3.1.2.3 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of 2-methyl-2-butanol, 0.06 ml of tributyrin and 20 mg of Savinase (protease from *Bacillus* sp.) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 22% ee was obtained in 39% yield (HPLC).
- 3.1.2.4 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of 2-methyl-2-butanol, 0.06 ml of tributyrin and 20 mg of subtilisin (protease from *Bacillus licheniformis*) at room temperature. After 4 days, (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 23% ee was obtained in 36% yield (HPLC).

09198427 112498
B5427 12486T60

3.1.3 Preparation of (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene using proteases

5 3.1.3.1 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of hexane, 0.06 ml of tributyrin and 120 U of Savinase (protease from *Bacillus* sp.) at room temperature. After 3-6 days, (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 44% ee was obtained in 46% yield (HPLC).

10 3.1.3.2 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of hexane, 0.06 ml of tributyrin and 20 mg of Alcalase (protease from *Bacillus licheniformis*) at room temperature. After 3-6 days, (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 44% ee was obtained in 35% yield (HPLC).

3.1.4 Preparation of (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene using lipases

20 3.1.4.1 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of 2-methyl-2-butanol, 0.03 ml of ethyl butyrate and 20 mg of Newlase F (lipase from *Rhizopus niveus*) at room temperature. After 1 week, (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 39% ee was obtained in 37% yield (HPLC).

30 3.1.4.2 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-2-cyclopentene were stirred with 1 ml of pyridine, 0.06 ml of tributyrin and 20 mg of lipase AK (lipase from *Pseudomonas fluorescens*) at room temperature. After 1 week, (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene with 30% ee was obtained in 10% yield (HPLC).

00198427.1.2498

3.1.4.3 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-
2-cyclopentene were stirred with 1 ml of
2-methyl-2-butanol, 0.06 ml of tributyrin and
20 mg of lipase AY (lipase from *Candida rugosa*)
at room temperature. After 1 week, (1S,4R)-1-
amino-4-(hydroxymethyl)-2-cyclopentene with 32%
ee was obtained in 13% yield (HPLC).

3.1.4.4 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-
2-cyclopentene were stirred with 1 ml of methyl
t-butyl ether, 0.06 ml of tributyrin and 20 mg
of lipase PS-CI (immobilized lipase from
Pseudomonas cepacia) at room temperature. After
1 week, (1S,4R)-1-amino-4-(hydroxymethyl)-2-
cyclopentene with 29% ee was obtained in 16%
yield (HPLC).

3.1.4.5 11 mg of racemic cis-1-amino-4-(hydroxymethyl)-
2-cyclopentene were stirred with 1 ml of methyl
t-butyl ether, 0.06 ml of tributyrin and 20 mg
of lipase PS (lipase from *Pseudomonas cepacia*)
at room temperature. After 1 week, (1S,4R)-1-
amino-4-(hydroxymethyl)-2-cyclopentene with 24%
ee was obtained in 22% yield (HPLC).

3.2 Racemate resolution using D-(-)-tartaric acid

3.2.1 A mixture of 8 g (70.6 mmol) of racemic 1-amino-
4-(hydroxymethyl)-2-cyclopentene and 10.6 g
(70.6 mmol) of D-(-)-tartaric acid in 186 g of
methanol were dissolved at the reflux tempera-
ture. The mixture was then cooled to 20°C over
2 h. At 43°C, seed crystals of the pure (1R,4S)-
1-amino-4-(hydroxymethyl)-2-cyclopentene D-
hydrogentartrate were added. The crystallized
product was filtered off and dried. Yield: 8.49 g
(45.6% based on racemic starting material) of
(1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene
D-hydrogentartrate, ee value: 91.1%. For purifi-

09198427 112498
B5427 12485T50

cation, 8.49 g (32.25 mmol) of the hydrogen-tartrate were suspended in 30 ml of methanol, and 2 equivalents of 30% sodium methoxide were added. The sodium tartrate was filtered off and the methanol was distilled off.

The residue was taken up in 35 ml of pentanol. Then, at 55°C, 1.5 g of HCl were introduced, and the solution was slowly cooled. At 40°C, the solution was seeded with (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene hydrochloride. 45 ml of acetone were then metered in, and the suspension was slowly cooled to 0°C and filtered, and the residue was dried. 3.91 g of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene hydrochloride having an ee value of >98% were obtained, corresponding to a yield, based on racemic (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene used, of 37%.

3.2.2 A mixture of 64 g of racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene (0.5 mol) and 75.2 g of D-(-)-tartaric acid in 1330 g of methanol was dissolved at the reflux temperature and then cooled to 20°C over 2 h. At 43°C, seed crystals of the pure 1R,4S-enantiomer were added. The crystallized product was filtered off and dried. Yield: 63.2 g (48.0% based on racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene) of (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene hydrogentartrate, ee value: 91.1%. The ee value in the mother liquor was 76.0%.

3.2.3 Recrystallization of 1R,4S-(4-amino-2-cyclopenten-1-yl)methanol D-hydrogentartrate

61.84 g of 1R,4S-(4-amino-2-cyclopenten-1-yl)-methanol D-hydrogentartrate (0.235 mol, ee value 91.1%) were dissolved in 752 g of methanol under reflux. The solution was cooled to 20°C within

09198427 112498

90 min, then the product was filtered off and washed with 64 g of cold methanol. Drying gave 54.56 g of 1R,4S-(4-amino-2-cyclopenten-1-yl)-methanol D-hydrogentartrate were obtained, ee value 99.4% (yield 88.2%, 42.3% based on racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene). This was used tel quel in the chloropurine synthesis.

3.2.4 Following the procedure of Example 3.2.2, but using 223 g of methanol and seeding at 50°C, the racemate was separated. The yield was 7.98 g (42.9% based on racemic (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene used).

3.3 Racemate resolution using L-(+)-tartaric acid

3.3.1 A mixture of 8 g (70.6 mmol) of racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene and 10.6 g (70.6 mmol) of L-(+)-tartaric acid in 186 g of methanol were dissolved at the reflux temperature. The mixture was then cooled to 20°C over 2 h. At 43°C, seed crystals of the pure (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene L-hydrogentartrate were added. The crystallized (1S,4R)-1-amino-4-(hydroxymethyl)-2-cyclopentene L-hydrogentartrate was filtered off and dried. (ee value: 91.1%). 14 g of 30% methanolic sodium methoxide were added to the mother liquor, then the methanol was evaporated. The residue was taken up in 35 ml of isobutanol, and the insoluble sodium tartrate was filtered off. At 55°C, 2 g of gaseous HCl were introduced into the filtrate. 38 ml of acetone were then added, and the mixture was left to cool to 10°C over the course of 1 h. After 1 h, the (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene hydrochloride was filtered off with suction and washed with 8 ml of acetone. Drying under reduced pressure gave the (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene

09198427 112493
B6427 2486T60

hydrochloride in a yield of 34 g, 31.6% based on racemic 1-amino-4-(hydroxymethyl)-2-cyclopentene with an ee value of >98%.

5 **Example 4: Preparation of (1R,4S)-amino-4-(hydroxymethyl)-2-cyclopentene hydrochloride**

4.1 Reduction of (-)-2-azabicyclo[2.2.1]hept-5-en-3-one

09498427 112498
06427 2486560
10 A 2 l autoclave (stainless steel type V4A) rendered inert with N₂, was charged with 61.4 g of sodium borohydride 97.5% (1.623 mol), 70.2 g of lithium chloride 98.5% (1.656 mol), 13.2 g of Celite and 1410 g of tetrahydrofuran. The autoclave was closed and heated to an internal temperature of 130°C and the contents stirred for 4.5 hours at this temperature (max. 8.0 bar).

15 After the autoclave had been cooled to about 60°C, the sodium salts insoluble in tetrahydrofuran (NaCl, NaBH₄) were filtered off. These were washed with 353 g of tetrahydrofuran, and the combined filtrates were reduced to about half in a stirred 1 l glass vessel by distillation at atmospheric pressure (distillate 1: about 710 g of tetrahydrofuran).
20 Further distillation, alternating with the portion-wise addition of a total of 936 g of dioxane then completed the solvent exchange (distillate 2: about 1289 g of tetrahydrofuran/dioxane).

25 The LiBH₄ suspension was cooled to about 60°C, and 56.7 g of (-)-2-azabicyclo[2.2.1]hept-5-en-3-one (97.5%) were added.

30 Starting at about 60°C, 132.5 g of methanol were metered in in exactly one hour at a rate such that a temperature range of 58-62°C was maintained. The mixture was then allowed to react for a further hour at 60°C. A further 397.0 g of methanol were then added (sample comprises an analytical yield of
35 70.5%), and the contents of the stirred vessel were cooled to 0°C. At this temperature, 90.0 g of HCl

09198427 112498
were introduced into the reaction mixture (slightly exothermic) and stirring was continued for a further hour at about 0°C. Distillation at atmospheric pressure (up to a head temperature of 75°C) removed the low-boiling fractions (methanol, borate) and about 70% of the dioxane (distillate 3: about 1093 g). Distillation under reduced pressure (about 30 mbar), alternating with the portionwise addition of a total of 282 g of 1-pentanol then completed the solvent exchange (distillate 4: about 240 g of dioxane/pentanol).

After a further 302 g of 1-pentanol had been added, the mixture was stirred for 1 hour at 50°C, and precipitated salts, about 39 g moist weight, were filtered off and washed with 200 g of 1-pentanol. The combined filtrates were reduced by redistillation under reduced pressure (about 20 mbar) (distillate 5: 235 g of 1-pentanol). Then, at about 50°C, 236 g of acetone were metered in, and the reaction mixture was seeded with a few crystals of (1R,4S)-amino-4-(hydroxymethyl)-2-cyclopentene. The mixture was cooled to 5°C over the course of 1 hour, and crystallization was completed by stirring the mixture for a further 6 h at 5°C.

The crystals were filtered off, washed with 63 g of acetone and dried at a max. 50°C in a vacuum drying cabinet (10 mbar). This gave 83.5 g of crude product* (content: 56.5%).

This corresponded to a yield of 61.4% based on (-)-2-azabicyclo[2.2.1]hept-5-en-3-one used.

4.2 Reduction of (±)-2-azabicyclo[2.2.1]hept-5-en-3-one

A 2 l autoclave (stainless steel type V4A) rendered inert with N₂, was charged with 41.56 g of sodium borohydride 97.5% (1.071 mol), 51.48 g of lithium chloride 98.5% (1.196 mol), 9.30 g of Celite and 955.0 g of tetrahydrofuran. The autoclave was closed

and heated to an internal temperature of 130°C and the contents stirred for 6 hours at this temperature (max. 6.3 bar).

5 After the autoclave had been cooled to about 60°C, the sodium salts insoluble in tetrahydrofuran (NaCl, NaBH₄) were filtered off. These were washed with 239.0 g of tetrahydrofuran, and the combined filtrates were reduced to about half in a stirred 1 l glass vessel by distillation at atmospheric pressure (distillate 1: about 590 g of THF). Further distillation, alternating with the portionwise addition of a total of 661.0 g of dioxane then completed the solvent exchange (distillate 2: about 685 g of tetrahydrofuran/dioxane).

15 The LiBH₄ suspension was cooled to about 60°C, and 36.0 g of 2-azabicyclo[2.2.1]hept-5-en-3-one (97.5%) were added.

Starting at about 60°C, 77.6 g of methanol were metered in in exactly one hour at a rate such that a temperature range of 58-62°C was maintained. The mixture was then allowed to react for a further hour at 60°C. A further 233.0 g of methanol were then added, and the contents of the stirred vessel were cooled to 0°C. At this temperature, 52.9 g of HCl were introduced into the reaction mixture (slightly exothermic) and stirring was continued for a further hour at about 0°C. Distillation at atmospheric pressure (up to a head temperature of 75°C) removed the low-boiling fractions (methanol, borate) and about 70% of the dioxane (distillate 3: about 700 g). Distillation under reduced pressure (about 30 mmol), alternating with the portionwise addition of a total of 169.4 g of 1-pentanol then completed the solvent exchange (distillate 4: about 183 g of dioxane/pentanol). After a further 127.1 g of 1-pentanol had been added, the mixture was stirred for 1 hour at 50°C, and precipitated salts, about 41 g moist weight, were filtered off and washed with 63.5 g of 1-pentanol. The combined filtrates were

09198427 112498

reduced by redistillation under reduced pressure (about 20 mbar) (distillate 5: 235 g of 1-pentanol). Then, at about 50°C, 238.0 g of acetone were metered in, and the reaction mixture was seeded with a few crystals of aminoalcohol hydrochloride salt. The mixture was cooled to 5°C over the course of one hour, and crystallization was completed by stirring the mixture for a further 6 hours at 5°C.

The crystals were filtered off, washed with 61.0 g of acetone and dried at a max. 50°C in a vacuum drying cabinet (10 mbar). This gave 50.0 g of crude product (content: about 50% of aminoalcohol hydrochloride salt).

This corresponded to a yield of 52.0% based on 2-azabicyclo[2.2.1]hept-5-en-3-one used.

Example 5: Preparation of acylated aminoalcohols

5.1 Preparation of (1R,4S)-N-BOC-1-amino-4-(hydroxymethyl)-2-cyclopentene (BOC = tert-butoxycarbonyl)

75 g of a solution of (1R,4S)-1-amino-4-hydroxymethyl-2-cyclopentene were adjusted to pH 8 using 30% strength NaOH, and 6 g of NaHCO₃ were added to the mixture. The mixture was heated to 52°C. Whilst stirring the mixture thoroughly, 60 ml of diisopropyl ether were added thereto and then, over the course of 2 h, a solution of 11.12 g of BOC anhydride in 18.2 ml of diisopropyl ether were metered in. The mixture was filtered over Celite, and the phases were separated. The aqueous phase was extracted with 65 ml of diisopropyl ether. The combined organic phases were washed with 45 ml of water, then evaporated to 37.5 g and heated to 50°C. 31 ml of n-hexane were added dropwise to the solution. After the mixture had been slowly cooled to 0°C (2 h), the title compound was filtered, washed with 12 ml of n-hexane/diisopropyl ether 1/1 and

09198427 112498

dried. This gave 6.75 g of product. The yield was 71%.

5.2 Preparation of (1R,4S)-N-acetyl-1-amino-4-(hydroxymethyl)-2-cyclopentene

5 25 g (1R,4S)-1-amino-4-(hydroxymethyl)-2-cyclopentene hydrochloride were dissolved in 182 ml of acetic anhydride, and at 0°C, a solution of 18.25 g of triethylamine in 60 ml of acetic anhydride were added thereto. The mixture was stirred at 80°C for 10 3 h, then cooled to room temperature. The triethylamine hydrochloride was filtered off and washed with 120 ml of n-hexane. The filtrate was evaporated. 300 ml of toluene were added to the residue, and the mixture was stirred at room temperature in the presence of 5.2 g of activated 15 carbon and 13 g of Celite for 20 min. The mixture was then filtered, and the filter cake was washed (3 x 40 ml of toluene), and the solvent was completely evaporated. 180 ml of methanol and 15.5 g of 20 K₂CO₃ were added to the residue, and the mixture was stirred at room temperature for 10 h. The suspension was filtered off and the filtrate evaporated. The residue was suspended in 750 ml of isopropyl acetate and boiled in the presence of 0.5 g of activated 25 carbon to reflux for 1.5 h. Following filtration of the activated carbon (70 - 80°C), the filtrate was cooled at 0°C overnight. The title compound was filtered, washed with 80 ml of cold isopropyl acetate and dried under reduced pressure to give 17.2 g 30 of product. The yield was 66%.

5.3 Preparation of (1R,4S)-N-butyryl-1-amino-4-(hydroxymethyl)-2-cyclopentene

34.7 g of (1R,4S)-1-amino-4-hydroxymethyl-2-cyclopentene hydrochloride and 2 g of N,N-4-dimethylaminopyridine were dissolved in 600 ml of methylene 35

09198427 112493 364211 2486160

chloride. The solution was cooled to 0°C. 52 g of triethylamine were then added dropwise (5 min). The mixture was stirred for a further 30 min. At 0°C, a solution of 35.2 g of butyryl chloride in 60 ml of methylene chloride was metered into the mixture over the course of 1 h. The mixture was stirred for a further 1.5 h at between 0 and 20°C, and then 600 ml of water were added thereto. Following phase separation, the aqueous phase was extracted with 600 ml of methylene chloride. The combined organic phases were washed 3 x 500 ml of 10% strength NaOH, then completely evaporated. The dried solid was dissolved in 120 ml of methanol. 5 g of K₂CO₃ were added to the solution, and the mixture was stirred for a further 2 h at room temperature. The inorganic salts were filtered off and washed with 20 ml of methanol. The filtrate was neutralized with 2N HCl. The suspension was filtered off, and the filter cake was washed with 20 ml of methanol. The filtrate was completely evaporated. The solid residue was dried and crystallized in 150 ml of toluene to give 28.5 g of the title compound. The yield was 67%.

Example 6: Preparation of [4(R)-(2-amino-6-chloropurine-9-yl)cyclopent-2-ene-1(S)-yl]methanol

6.1 Preparation of [4(R)-(2-amino-6-chloropurine-9-yl)-cyclopent-2-ene-1(S)-yl]methanol starting from 1R,4S-(4-amino-2-cyclopenten-1-yl)methanol D-hydrogentartrate

47.4 g of 1R,4S-(4-amino-2-cyclopenten-1-yl)methanol D-hydrogentartrate (0.18 mol, ee >98%) in 200 ml of ethanol were introduced initially. At room temperature, 54.6 g of NaHCO₃ (0.65 mol) and 37.3 g (0.18 mol) of N-(2-amino-4,6-dichloro-4-pyrimidyl)-formamide were added, boiled for 9 h under reflux and then cooled to room temperature. The salts were

09198427 112493

filtered off and then washed with 50 ml of ethanol. The filtrate was concentrated to 280 g on a rotary evaporator. 18.4 g of HCl gas were introduced into the resulting solution at $T < 25^{\circ}\text{C}$, then 95.5 g (0.9 mol) of trimethyl orthoformate were added, and the whole was heated to 40°C (10 min). At this temperature, the mixture was seeded with chloropurine hydrochloride. After 2 h at 42°C , the product crystallized out. The suspension was cooled to 15°C . The product was filtered and then washed with 3×50 ml of ethanol, then dried at 50°C under reduced pressure. The yield was 41.9 g (75.8%). Beige powder, content (HPLC): 95.0%.

6.2 Preparation of [4(R)-(2-amino-6-chloropurine-9-yl)-cyclopent-2-ene-1(S)-yl]methanol starting from (-)-2-azabicyclo[2.2.1]hept-5-en-3-one

A 2 l autoclave (stainless steel type V4A) rendered inert with N_2 , was charged with 61.4 g of sodium borohydride 97.5% (1.623 mol), 70.2 g of lithium chloride 98.5% (1.656 mol), 13.2 g of Celite and 1410 g of tetrahydrofuran. The autoclave was closed and heated to an internal temperature of 130°C and the contents stirred for 4.5 hours at this temperature (max. 8.0 bar). After the autoclave had been cooled to about 60°C , the sodium salts insoluble in tetrahydrofuran (NaCl , NaBH_4) were filtered off. These were washed with 353 g of tetrahydrofuran, and the combined filtrates were reduced to about half in a stirred 1 l glass vessel by distillation at atmospheric pressure (distillate 1: about 710 g of tetrahydrofuran). Further distillation, alternating with the portionwise addition of a total of 936 g of dioxane then completed the solvent exchange (distillate 2: about 1289 g of tetrahydrofuran/dioxane).

The LiBH_4 suspension was cooled to about 60°C , and 56.7 g of (-)-2-azabicyclo[2.2.1]hept-5-en-3-one

00198427 112493

(97.5%/0.507 mol) were added.

Starting at about 60°C, 132.5 g of methanol were metered in in exactly one hour at a rate such that a temperature range of 58-62°C was maintained. The mixture was then allowed to react for a further hour at 60°C. A further 397.0 g of methanol were then added (sample comprises an analytical yield of 70.5%), and the contents of the stirred vessel were cooled to 0°C. At this temperature, 90.0 g of HCl were introduced into the reaction mixture (slightly exothermic) and stirring was continued for a further hour at about 0°C. The solution was evaporated on a rotary evaporator at 50°C under reduced pressure, 200 ml of methanol were added and the methanol was removed again (filtration with suction of the methyl borate). The procedure was repeated using a further 200 ml of methanol. 250 ml of ethanol were added to the oil obtained (253.4 g comprise 3.16% of amino-alcohol; this corresponded to 0.360 mol), and the mixture was poured into a 1 l double-jacketed stirred vessel. At room temperature, 72.6 g of NaHCO₃ (0.86 mol) and 74.6 g (0.360 mol) of N-(2-amino-4,6-dichloro-4-pyrimidyl)formamide were added, the mixture was refluxed for 9 h and cooled to room temperature, and the salts were filtered off and then washed with 100 ml of ethanol. The filtrate on the rotary evaporator was concentrated to 560 g. 63.4 g of HCl gas were introduced into the resulting solution at T<25°C, then 191.0 g (1.80 mol) of trimethyl orthoformate were added, and the mixture was heated to 40°C (10 min). At this temperature, the mixture was seeded with chloropurine hydrochloride, and left to crystallize for 2 h at 42°C. The suspension was cooled to 15°C. The product was filtered and then washed with 3 x 50 ml of ethanol, then dried at 50°C under reduced pressure. The yield was 66.0 g (59.7%). Beige powder, content (HPLC): 89.3%. This corresponded to a yield of 42.4% based on the Vince lactam used.

09198427-112498